Thurston County

Youth & Tobacco Data Summary 10th Grade

| Vear | 2000 | |
|-------|------|--|
| ı caı | 2000 | |

| 100.0 % 100. | | | 1001 2000 | County % | State % | |
|--|---|-------------------------|---------------------------------|--------------------------|---------------|--|
| Female | 10th Grade S | Students Surveyed | n= 1,932 | 100.0 % | 100.0 % | |
| Male 46.7 % 48.4 % Race/Ethnicity 22.8 % 24.2 % White 22.8 % 24.2 % White 22.8 % 24.2 % Name County %+/- Margin of Error Margin of Error Past 30 days, youth who have smoked cigarettes at least once Female n = 1018 21.8 %+/- 2.5 21.8 %+/- 1.5 21.8 %+/- 1.2 21.8 %+/- 1.2 21.8 %+/- 1.2 21.8 %+/- 1.2 <th< td=""><td>Gender</td><td></td><td></td><td></td><td></td></th<> | Gender | | | | | |
| Students of Color Ywhite 12.8 % 24.2 % 77.2 % 75.8 % 75.8 % 77.2 % 75.8 % 75.8 % 77.2 % 75.8 % | Female | | | 53.3 % | 51.6 % | |
| Students of Color | Male | | | 46.7 % | 48.4 % | |
| White 77.2 % 75.8 % County %+/- Margin of Error State %+/- Margin of Error Past 30 days, youth who have smoked cigarettes at least once 21.8 %+/- 2.5 21.8 %+/- 2.5 Female n= 1018 21.8 %+/- 2.5 21.8 %+/- 2.2 Male n= 890 20.4 %+/- 2.6 17.5 %+/- 2.1 Total n= 1908 3.9 %+/- 1.2 4.8 %+/- 1.5 Past 30 days, youth who have smoked cigars at least once Female n= 1016 3.9 %+/- 1.2 4.8 %+/- 1.2 1.1 %+/- 1.2 1.2 %+/- 1.7 1.1 %+/- 2.1 11.2 %+/- 1.7 1.1 %+/- 2.1 11.2 %+/- 1.0 1.2 %+/- 1.0 1.2 %+/- 1.0 1.2 %+/- 1.0 1.0 %+/- 1.0 1.2 %+/- 1.0 1.0 %+/- 0.6 1.2 %+/- 0.5 1.2 %+/- 0.6 1.2 %+/- 0.6 1.2 | Race/Ethnici | ty | | | | |
| County %+/- Margin of Error | Students of | Color | | 22.8 % | 24.2 % | |
| Margin of Error Margin of Error Margin of Error | White | | | 77.2 % | 75.8 % | |
| Past 30 days, youth who have smoked cigarettes at least once Female n = 1018 21.8 %+/- 2.5 21.8 %+/- 2.1 17.5 %+/- 2.1 1908 21.2 %+/- 1.8 19.8 %+/- 1.5 19 | | | | County %+/- | State %+/- | |
| Past 30 days, youth who have smoked cigarettes at least once Female | | | | Margin | Margin | |
| ## least once Female | | | | _ | _ | |
| Female | | | ked cigarettes | | | |
| Total n= 1908 | | | | 21.8 %+/- 2.5 | 21.8 %+/- 2.2 | |
| Past 30 days, youth who have smoked cigars at least once Female n= 1016 3.9 %+/- 1.2 4.8 %+/- 1.2 Male n= 891 11.6 %+/- 2.1 11.2 %+/- 1.7 Total n= 1907 7.6 %+/- 1.2 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 1013 1.1 %+/- 0.6 1.2 %+/- 0.8 Male n= 886 2.7 %+/- 1.1 2.7 %+/- 0.8 Total n= 1899 1.9 %+/- 0.6 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 1.0 4.6 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 2.3.6 %+/- 2.6 23.8 %+/- 0.9 Male n= 877 2.4 8 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 2.4 8 %+/- 2.4 | Male | n= 890 | | 20.4 %+/- 2.6 | 17.5 %+/- 2.1 | |
| Female n= 1016 3.9 %+/- 1.2 4.8 %+/- 1.2 Male n= 891 11.6 %+/- 2.1 11.2 %+/- 1.7 Total n= 1907 7.6 %+/- 1.2 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 1013 1.1 %+/- 0.6 1.2 %+/- 0.6 Male n= 886 2.7 %+/- 1.1 2.7 %+/- 0.8 Total n= 1899 1.9 %+/- 0.6 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 <td colspan<="" td=""><td>Total</td><td>n= 1908</td><td></td><td></td><td></td></td> | <td>Total</td> <td>n= 1908</td> <td></td> <td></td> <td></td> | Total | n= 1908 | | | |
| Female n= 1016 3.9 %+/- 1.2 4.8 %+/- 1.2 Male n= 891 11.6 %+/- 2.1 11.2 %+/- 1.7 Total n= 1907 7.6 %+/- 1.2 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 1013 1.1 %+/- 0.6 1.2 %+/- 0.6 Male n= 886 2.7 %+/- 1.1 2.7 %+/- 0.8 Total n= 1899 1.9 %+/- 0.6 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 <td colspan<="" td=""><td>Past 30 days</td><td>, youth who have smo</td><td>ked cigars at least once</td><td></td><td></td></td> | <td>Past 30 days</td> <td>, youth who have smo</td> <td>ked cigars at least once</td> <td></td> <td></td> | Past 30 days | , youth who have smo | ked cigars at least once | | |
| Male n= 891 11.6 %+/- 2.1 11.2 %+/- 1.7 Total n= 1907 7.6 %+/- 1.2 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 1013 1.1 %+/- 0.6 1.2 %+/- 0.6 Male n= 886 2.7 %+/- 1.1 2.7 %+/- 0.8 Total n= 1899 1.9 %+/- 0.6 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | • | | • | 3.9 %+/- 1.2 | 4.8 %+/- 1.2 | |
| Total n= 1907 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 1013 Male n= 886 Total n= 1899 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 Male n= 884 Total n= 1893 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 878 Total n= 1870 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 878 Total n= 1870 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Past 30 days, youth who have used any tobacco product at least once | | | | | 11.2 %+/- 1.7 | |
| at least once Female n= 1013 | | | | | | |
| Female n= 1013 1.1 %+/- 0.6 1.2 %+/- 0.8 Male n= 886 2.7 %+/- 1.1 2.7 %+/- 0.8 Total n= 1899 1.9 %+/- 0.6 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Past 30 days | , youth who have smo | ked tobacco in a pipe | | | |
| Male n= 886 2.7 %+/- 1.1 2.7 %+/- 0.8 Total n= 1899 1.9 %+/- 0.6 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | at least once | e | | | | |
| Total n= 1899 1.9 %+/- 0.6 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 Male n= 884 Total n= 1893 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 Male n= 878 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 878 Past 30 days, youth who have used any tobacco product at least once Female n= 878 Past 30 days, youth who have used any tobacco product at least once Female n= 992 Male n= 877 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 | Female | n= 1013 | | 1.1 %+/- 0.6 | 1.2 %+/- 0.6 | |
| Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 1009 | Male | n= 886 | | 2.7 %+/- 1.1 | 2.7 %+/- 0.8 | |
| at least once Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Total | n= 1899 | | 1.9 %+/- 0.6 | 1.9 %+/- 0.5 | |
| Female n= 1009 4.5 %+/- 1.3 4.5 %+/- 1.1 Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Past 30 days | , youth who have smo | ked bidis (flavored cigarettes) | | | |
| Male n= 884 5.7 %+/- 1.5 4.7 %+/- 1.2 Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | at least once | e | | | | |
| Total n= 1893 5.1 %+/- 1.0 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Female | n= 1009 | | 4.5 %+/- 1.3 | 4.5 %+/- 1.1 | |
| Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Male | n= 884 | | 5.7 %+/- 1.5 | 4.7 %+/- 1.2 | |
| (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Total | n= 1893 | | 5.1 %+/- 1.0 | 4.6 %+/- 0.8 | |
| (chewing tobacco, snuff or dip) at least once Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Past 30 days | , youth who have used | l smokeless tobacco | | | |
| Female n= 992 0.8 %+/- 0.6 1.4 %+/- 0.8 Male n= 878 9.2 %+/- 1.9 7.9 %+/- 1.6 Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | (chewing to | bacco, snuff or dip) at | least once | | | |
| Male Total n= 878 n= 1870 9.2 %+/- 1.9 | | | | 0.8 %+/- 0.6 | 1.4 %+/- 0.8 | |
| Total n= 1870 4.8 %+/- 1.0 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Male | n= 878 | | 9.2 %+/- 1.9 | 7.9 %+/- 1.6 | |
| at least once Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | Total | | | | | |
| Female n= 992 23.6 %+/- 2.6 23.8 %+/- 2.3 Male n= 877 26.7 %+/- 2.9 24.8 %+/- 2.4 | • | | l any tobacco product | | | |
| Male n= 877 <u>26.7 %+/- 2.9</u> 24.8 %+/- 2.4 | at least onc | e | | | | |
| | Female | n= 992 | | 23.6 %+/- 2.6 | | |
| Total n= 1869 25.1 %+/- 2.0 24.4 %+/- 1.6 | Male | n= 877 | | <u>26.7</u> %+/- 2.9 | | |
| | Total | n= 1869 | | 25.1 %+/- 2.0 | 24.4 %+/- 1.6 | |

Source: Washington State Department of Health Tobacco Prevention and Control Assessment and Evaluation n = number of survey respondents for county.

* = Rate not calculated because number of responses was less than 15.

Thurston County

Youth & Tobacco Data Summary 10th Grade

| Year | 20 | OO. |
|------|----|-----|
| | | |

| Teal 2000 | County %+/- Margin of Error | State %+/- Margin of Error |
|--|--|---|
| About how many cigarettes have you smoked in your entire life? None 1/2 pack or less 1 - 4 packs 5 or more packs | n= 1877 49.8 %+/- 2.3 23.1 %+/- 1.9 8.5 %+/- 1.3 18.6 %+/- 1.8 | 49.9 %+/- 1.9 24.7 %+/- 1.6 9.3 %+/- 1.1 16.1 %+/- 1.4 |
| Do you want to stop using tobacco right now? (among current tobacco users) Yes | n= 461 59.4 %+/- 4.5 | 56.8 %+/- 4.0 |
| Have you ever participated in a program to help you quit using tobacco? (among current tobacco users) Yes | n= 604 5.0 %+/- 1.7 | 6.7 %+/- 1.7 |
| During past 12 months, did you ever try to quit using tobacco? (cigarettes, cigars, or chew/dip)(among current tobacco users) Yes | n= 353 59.2 %+/- 5.1 | 58.3 %+/- 4.9 |
| Do you think you will smoke a cigarette at any time during the next year? Definitely no | n= 1915 57.6 %+/- 2.2 | 56.2 %+/- 1.8 |
| If one of your best friends offered you a cigarette, would you smoke it? Definitely no | n= 1916 61.3 %+/- 2.2 | 59.1 %+/- 1.8 |
| Susceptibility Index (combination of the previous two questions, either of which receives any answer other than 'Definitely no') Susceptible | n= 1916 44.4 %+/- 2.2 | 46.8 %+/- 1.8 |
| If one of your best friends offered you a cigarette, would you smoke it? (from among youth who did not smoke in last 30 days) Definitely no | n= 1507 76.6 %+/- 2.1 | 72.8 %+/- 1.8 |
| Do you think the smoke from other people's cigarettes (second-hand smoke) is harmful to you? Definitely yes or Probably yes | n= 1912 92.4 %+/- 1.2 | 92.2 %+/- 1.0 |
| Do you think it is safe to smoke for only a year or two, as long as you quit after that? Definitely no | n= 1914 67.9 %+/- 2.1 | 67.7 %+/- 1.7 |

Thurston County

Youth & Tobacco Data Summary 10th Grade Year 2000

| | County %+/- Margin of Error | State %+/- Margin of Error |
|---|-----------------------------------|----------------------------------|
| Would you ever use or wear something that has a tobacco company name or picture on it such as a lighter, t-shirt, hat, or sunglasses? Definitely no | n= 1915 39.6 %+/- 2.2 | 39.7 %+/- 1.8 |
| During last 12 months, did you buy or receive anything that has a tobacco name or picture on it? Yes | n= 1918 20.0 %+/- 1.8 | 18.7 %+/- 1.4 |
| During past 7 days, on how many days were you in the same room with someone who was smoking cigarettes? 3 or more days | n= 766 34.2 %+/- 3.4 | 28.7 %+/- 2.7 |
| During past 7 days, on how many days did you ride in a car with someone who was smoking cigarettes? 3 or more days | n= 764 25.9 %+/- 3.1 | 24.6 %+/- 2.5 |
| During this school year, did you practice ways to say 'NO' to tobacco in any of your classes (for example, by role playing)? Yes | n= 1913 28.6 %+/- 2.0 | 27.5 %+/- 1.6 |
| During past 30 days, have you seen or heard commercials on TV, the Internet, or on the radio about the dangers of cigarette smoking? More than once a week | n= 1911 81.6 %+/- 1.7 | 77.6 %+/- 1.6 |
| How wrong would most adults in your neighborhood think it was for kids your age to smoke cigarettes? Very wrong | n= 1913 36.9 %+/- 2.2 | 39.4 %+/- 1.8 |
| If you wanted to get some tobacco, how easy would it be for you to get some? Very hard or Sort of hard | n= 1902 27.4 %+/- 2.0 | 27.8 %+/- 1.6 |
| Has either of your parents (or guardians) discussed the dangers of tobacco use with you? Yes | n= 1917 67.7 %+/- 2.1 | 65.8 %+/- 1.7 |
| Does anyone who lives with you now smoke cigarettes? Yes | n= 767 41.3 %+/- 3.5 | 37.4 %+/- 2.9 |